



14 wks cold; in grnhse 21 Jan. (9513)



19 wks cold; in grnhse 18 Mar. (9707)



15 wks cold; in grnhse 4 Feb. (9619)



21 wks cold; in grnhse 1 Apr. (9979)



16 wks cold; in grnhse 18 Feb. (9808)



23 wks cold; in grnhse 15 Apr. (9959)



17 wks cold; in grnhse 4 Mar (9265)

Effect of cold-weeks and Bonzi drenches on appearance of 6" 'Agrass White' tulips. Cold weeks vary between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Bonzi applied as a soil drench 1-2 days after placing in the greenhouse. 2002.





12 wks cold; in grnhse 15 Dec. (1220)



16 wks cold; in grnhse 23 Feb. (0755)



14 wks cold; in grnhse 5 Jan. (1319)



18 wks cold; in grnhse 8 Mar. (1060)



15 wks cold; in grnhse 19 Jan. (0076)



20 wks cold; in grnhse 22 Mar. (1289)



17 wks cold; in grnhse 2 Feb (0373)



22 wks cold; in grnhse 12 Apr. (1498)

Effect of cold-weeks and Bonzi drenches on appearance of 6" 'Agrass White' tulips. Cold weeks vary between panels from 12-24 weeks. In each panel, L to R: Control, 1 and 2 mg Bonzi applied as a soil drench 1-2 days after placing in the greenhouse. 2004.





12 wks cold; in grnhse 20 Dec. (9466)



17 wks cold; in grnhse 28 Feb. (9921)




14 wks cold; in grnhse 10 Jan. (9586)

N/A 19 wks cold; in grnhse 14 Mar. (xxx)



15 wks cold; in grnhse 25 Jan. (9589)





16 wks cold; in grnhse 14 Feb (9788)



23 wks cold; in grnhse 11 Apr. (9649) Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Agrass White'

tulips. Cold duration varies between panels from 12-24 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo applied as a soil drench 1-3 days after placing in the greenhouse.



20 wks cold; in grnhse 29 Mar. (2920)



n/a 16 wks cold; in grnhse 25 Jan. (xxx)

22 wks cold; in grnhse 12 Apr. (2986)



17 wks cold; in grnhse 1 Mar. (2507)



18 wks cold; in grnhse 15 Mar. (2736)

Effect of cold-weeks and PGR drenches on appearance of 6" tulip. Cold duration varies between panels from 16-22 weeks. In each panel, L to R: Control, 1 and 2 mg Bonzi, 0.5 and 1 mg/pot Topflor applied as a soil drench (2-4 days after placing in the greenhouse). 2011.