

Bright Parrot 2004

N/A

12 wks cold; in grnhse 15 Dec. (xxx)



16 wks cold; in grnhse 23 Feb. (0742)



14 wks cold; in grnhse 5 Jan. (1266)



18 wks cold; in grnhse 8 Mar. (1031)



15 wks cold; in grnhse 19 Jan. (0011)



20 wks cold; in grnhse 22 Mar. (1294)



17 wks cold; in grnhse 2 Feb (0257)



22 wks cold; in grnhse 12 Apr. (1495)

Effect of cold-weeks and Bonzi drenches on appearance of 6" 'Bright Parrot' tulips. Cold weeks vary between panels from 12-24 weeks. In each panel, L to R: Control, 1 and 2 mg Bonzi applied as a soil drench 1-2 days after placing in the greenhouse. 2004.

Bright Parrot 2005



12 wks cold; in grnhse 20 Dec. (9325)



17 wks cold; in grnhse 28 Feb. (9866)



14 wks cold; in grnhse 10 Jan. (9492)

N/A

19 wks cold; in grnhse 14 Mar. (xxx)



15 wks cold; in grnhse 25 Jan. (9587)

N/A

21 wks cold; in grnhse 28 Mar. (xxx)



16 wks cold; in grnhse 14 Feb (9791)



23 wks cold; in grnhse 11 Apr. (9644)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Bright Parrot' tulips. Cold duration varies between panels from 12-24 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo applied as a soil drench 1-3 days after placing in the greenhouse.

Bright Parrot 2006



13 wks cold; in grnhse 26 Dec. (0341)



17 wks cold; in grnhse 27 Feb. (2136)



14 wks cold; in grnhse 9 Jan. (0559)



19 wks cold; in grnhse 13 Mar. (2415)



15 wks cold; in grnhse 23 Jan. (0687)

N/A
21 wks cold; in grnhse 27 Mar. (xxx)



16 wks cold; in grnhse 13 Feb (1751)



23 wks cold; in grnhse 10 Apr. (2984)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Bright Parrot' tulips. Cold duration varies between panels from 13-24 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo applied as a soil drench 1-3 days after placing in the greenhouse. Experiment 2006-T1.

Bright Parrot 2007



14 wks cold; in grnhse 2 Jan. (0038)



17 wks cold; in grnhse 26 Feb. (0342)



15 wks cold; in grnhse 15 Jan. (0085)



19 wks cold; in grnhse 12 Mar. (0404)



16 wks cold; in grnhse 29 Jan. (0200)



21 wks cold; in grnhse 26 Mar. (1316)



17 wks cold; in grnhse 12 Feb (0232)

N/A

23 wks cold; in grnhse 16 Apr. (xxx)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Bright Parrot' tulips. Cold duration varies between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo applied as a soil drench 4-6 days after placing in the greenhouse.

Bright Parrot 2009



14 wks cold; in grnhse 6 Jan. (1050)



19 wks cold; in grnhse 10 Mar. (2016)



16 wks cold; in grnhse 20 Jan. (1200)



21 wks cold; in grnhse 24 Mar. (2079)



17 wks cold; in grnhse 3 Feb Jan. (1376)

n/a

23 wks cold; in grnhse 14 April (xxx)



18 wks cold; in grnhse 24 Feb. (1508)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6” ‘Bright Parrot’ tulips. Cold duration varies between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo, 0.5 and 1 mg/pot Topflor applied as a soil drench (2-7 days after placing in the greenhouse).