

Freeman 2009



14 wks cold; in grnhse 6 Jan. (1100)



19 wks cold; in grnhse 10 Mar. (2029)



16 wks cold; in grnhse 20 Jan. (1269)



21 wks cold; in grnhse 24 Mar. (2085)



17 wks cold; in grnhse 3 Feb Jan. (1412)

n/a

23 wks cold; in grnhse 14 April (xxx)



18 wks cold; in grnhse 24 Feb. (1590)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Freeman' tulips. Cold duration varies between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo, 0.5 and 1 mg/pot Topflor applied as a soil drench (2-7 days after placing in the greenhouse).

**Freeman (11/12 cm) 2010**



14 wks cold; in grnhse 4 Jan. (0105)



19 wks cold; in grnhse 12 Mar. (0706)



16 wks cold; in grnhse 21 Jan. (0309)



17 wks cold; in grnhse 5 Feb. (0457)



18 wks cold; in grnhse 1 Mar. (0522)

Effect of cold-weeks and paclobutrazol (Bonzi) drenches on appearance of 6'' 'Freeman' tulips. Cold duration varies between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo, 0.5 and 1 mg/pot Topflor applied as a soil drench (2-7 days after placing in the greenhouse).