

Frohnleiten 2009



14 wks cold; in grnhse 6 Jan. (1095)



19 wks cold; in grnhse 10 Mar. (2025)



16 wks cold; in grnhse 20 Jan. (1276)



21 wks cold; in grnhse 24 Mar. (2092)



17 wks cold; in grnhse 3 Feb Jan. (1415)

n/a

23 wks cold; in grnhse 14 April (xxx)



18 wks cold; in grnhse 24 Feb. (1587)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Frohnleiten' tulips. Cold duration varies between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo, 0.5 and 1 mg/pot Topflor applied as a soil drench (2-7 days after placing in the greenhouse).

Frohnleiten 2010



14 wks cold; in grnhse 4 Jan. (0098)



19 wks cold; in grnhse 12 Mar. (0709)



16 wks cold; in grnhse 21 Jan. (0315)



17 wks cold; in grnhse 5 Feb. (0446)

n/a

18 wks cold; in grnhse 1 Mar. (xxx)

Effect of cold-weeks and paclobutrazol (Bonzi) drenches on appearance of 6" 'Frohnleiten' tulips. Cold duration varies between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo, 0.5 and 1 mg/pot Topflor applied as a soil drench (2-7 days after placing in the greenhouse).