

Herman Emmink 2006

N/A

13 wks cold; in grnhse 26 Dec. (xxx)



14 wks cold; in grnhse 9 Jan. (0769)



15 wks cold; in grnhse 23 Jan. (1123)



16 wks cold; in grnhse 13 Feb (1951)

N/A

17 wks cold; in grnhse 27 Feb. (xxx)

N/A

19 wks cold; in grnhse 13 Mar. (xxx)

N/A

21 wks cold; in grnhse 27 Mar. (xxx)

N/A

23 wks cold; in grnhse 10 Apr. (xxx)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Herman Emmink' tulips. Cold duration varies between panels from 13-24 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo applied as a soil drench 1-3 days after placing in the greenhouse. Experiment 2006-T1.

Herman Emmink 2008



14.5 wks cold; in grnhse 7 Jan. (0065)



19 wks cold; in grnhse 11 Mar. (0715)



16 wks cold; in grnhse 22 Jan. (011)

N/A

21 wks cold; in grnhse 25 Mar. (xxx)



17 wks cold; in grnhse 5 Feb Jan. (0454)



23 wks cold; in grnhse 16 Mar. (0797)

N/A

18 wks cold; in grnhse 26 Feb. (xxx)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6” ‘Herman Emmink’ tulips. Cold duration varies between panels from 14.5-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo applied as a soil drench (2-7 days after placing in the greenhouse, depending on growth).