

**Kikomachi 2002**



14 wks cold; in grnhse 21 Jan. (9509)



19 wks cold; in grnhse 18 Mar. (9707)



15 wks cold; in grnhse 4 Feb. (9621)



21 wks cold; in grnhse 1 Apr. (9989)



16 wks cold; in grnhse 18 Feb. (9876)



23 wks cold; in grnhse 15 Apr. (9964)



17 wks cold; in grnhse 4 Mar (9262)

Effect of cold-weeks and Bonzi drenches on appearance of 6" 'Kikomachi' tulips. Cold weeks vary between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Bonzi applied as a soil drench 1-2 days after placing in the greenhouse. 2002.

Kikomachi 2003



12 wks cold; in grnhse 23 Dec. (3391)



17 wks cold; in grnhse 3 Mar. (3732)



14 wks cold; in grnhse 13 Jan. (3444)



19 wks cold; in grnhse 17 Mar. (xxx)



16 wks cold; in grnhse 27 Jan. (3513)



21 wks cold; in grnhse 31 Mar. (4797)



16 wks cold; in grnhse 17 Feb(3702)



24 wks cold; in grnhse 21 Apr. (6209)

Effect of cold-weeks and Bonzi drenches on appearance of 6" 'Kikomachi' tulips. Cold weeks vary between panels from 12-24 weeks. In each panel, L to R: Control, 1 and 2 mg Bonzi applied as a soil drench 1-2 days after placing in the greenhouse. 2003.

Kikomachi 2004



12 wks cold; in grnhse 15 Dec. (0027)



16 wks cold; in grnhse 23 Feb. (0738)



14 wks cold; in grnhse 5 Jan. (1516)



18 wks cold; in grnhse 8 Mar. (1014)



15 wks cold; in grnhse 19 Jan. (0122)



20 wks cold; in grnhse 22 Mar. (1253)



17 wks cold; in grnhse 2 Feb (0328)



22 wks cold; in grnhse 12 Apr. (1483)

Effect of cold-weeks and Bonzi drenches on appearance of 6" 'Kikomachi' tulips. Cold weeks vary between panels from 12-24 weeks. In each panel, L to R: Control, 1 and 2 mg Bonzi applied as a soil drench 1-2 days after placing in the greenhouse. 2004.

**Kikomachi 2009**



14 wks cold; in grnhse 6 Jan. (1167)



19 wks cold; in grnhse 10 Mar. (1859)



16 wks cold; in grnhse 20 Jan. (1266)



21 wks cold; in grnhse 24 Mar. (2052)



17 wks cold; in grnhse 3 Feb Jan. (1419)

n/a

23 wks cold; in grnhse 14 April (xxx)



18 wks cold; in grnhse 24 Feb. (1524)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Kikomachi' tulips. Cold duration varies between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo, 0.5 and 1 mg/pot Topflor applied as a soil drench (2-7 days after placing in the greenhouse).

## Kikomachi 2010



14 wks cold; in grnhse 4 Jan. (0111)



19 wks cold; in grnhse 12 Mar. (0672)



16 wks cold; in grnhse 21 Jan. (0275)



17 wks cold; in grnhse 5 Feb. (0414)



18 wks cold; in grnhse 1 Mar. (0507)

Effect of cold-weeks and paclobutrazol (Bonzi) drenches on appearance of 6" 'Kikomachi' tulips. Cold duration varies between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo, 0.5 and 1 mg/pot Topflor applied as a soil drench (2-7 days after placing in the greenhouse).