

King's Cloak 2003



12 wks cold; in grnhse 23 Dec. (3392)




14 wks cold; in grnhse 13 Jan. (3446)



16 wks cold; in grnhse 27 Jan. (3515)







17 wks cold; in grnhse 3 Mar. (3766)



19 wks cold; in grnhse 17 Mar. (4024)



21 wks cold; in grnhse 31 Mar. (4795)





King's Cloak 2004



12 wks cold; in grnhse 15 Dec. (1104)



14 wks cold; in grnhse 5 Jan. (1527)



15 wks cold; in grnhse 19 Jan. (0123)



17 wks cold; in grnhse 2 Feb (0377)22 wks cold; in grnhse 12 Apr. (1502)Effect of cold-weeks and Bonzi drenches on appearance of 6" 'King's Cloak' tulips. Coldweeks vary between panels from 12-24 weeks. In each panel, L to R: Control, 1 and 2 mgBonzi applied as a soil drench 1-2 days after placing in the greenhouse. 2004.



16 wks cold; in grnhse 23 Feb. (0761)



18 wks cold; in grnhse 8 Mar. (1061)



20 wks cold; in grnhse 22 Mar. (1318)





King's Cloak 2005



12 wks cold; in grnhse 20 Dec. (9467)



14 wks cold; in grnhse 10 Jan. (9559)



15 wks cold; in grnhse 25 Jan. (9591)



16 wks cold; in grnhse 14 Feb (9792)



17 wks cold; in grnhse 28 Feb. (9909)

N/A 19 wks cold; in grnhse 14 Mar. (xxx)

N/A 21 wks cold; in grnhse 28 Mar. (xxx)



23 wks cold; in grnhse 11 Apr. (9646) Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'King's Cloak' tulips. Cold duration varies between panels from 12-24 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo applied as a soil drench 1-3 days after placing in the greenhouse.