

Mary Belle 2006



13 wks cold; in grnhse 26 Dec. (0283)



17 wks cold; in grnhse 27 Feb. (2106)



14 wks cold; in grnhse 9 Jan. (0388)




19 wks cold; in grnhse 13 Mar. (2376)



15 wks cold; in grnhse 23 Jan. (0682)

16 wks cold; in grnhse 13 Feb (1748)





23 wks cold; in grnhse 10 Apr. (2978)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Mary Belle' tulips. Cold duration varies between panels from 13-24 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo applied as a soil drench 1-3 days after placing in the greenhouse. Experiment 2006-T1.



Mary Belle 2007



14 wks cold; in grnhse 2 Jan. (0039)



17 wks cold; in grnhse 26 Feb. (0338)



15 wks cold; in grnhse 15 Jan. (0078)



19 wks cold; in grnhse 12 Mar. (0407)



16 wks cold; in grnhse 29 Jan. (0166)



21 wks cold; in grnhse 26 Mar. (1313)



17 wks cold; in grnhse 12 Feb (0230)



23 wks cold; in grnhse 16 Apr. (1835) Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Mary Belle' tulips. Cold duration varies between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo applied as a soil drench 4-6 days after placing in the greenhouse.