

Toplips 2008



14.5 wks cold; in grnhse 7 Jan. (0053)

N/A

19 wks cold; in grnhse 11 Mar. (xxx)



16 wks cold; in grnhse 22 Jan. (0173)

N/A

21 wks cold; in grnhse 25 Mar. (xxx)



17 wks cold; in grnhse 5 Feb Jan. (0448)



23 wks cold; in grnhse 16 Mar. (0791)

N/A

18 wks cold; in grnhse 26 Feb. (xxx)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6" 'Top Lips' tulips. Cold duration varies between panels from 14.5-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo applied as a soil drench (2-7 days after placing in the greenhouse, depending on growth).

Toplips 2009



14 wks cold; in grnhse 6 Jan. (1223)



19 wks cold; in grnhse 10 Mar. (2032)



16 wks cold; in grnhse 20 Jan. (1326)



21 wks cold; in grnhse 24 Mar. (2098)



17 wks cold; in grnhse 3 Feb Jan. (1461)

n/a

23 wks cold; in grnhse 14 April (xxx)



18 wks cold; in grnhse 24 Feb. (1633)

Effect of cold-weeks and paclobutrazol (Piccolo) drenches on appearance of 6” ‘Toplips’ tulips. Cold duration varies between panels from 14-23 weeks. In each panel, L to R: Control, 1 and 2 mg Piccolo, 0.5 and 1 mg/pot Topflor applied as a soil drench (2-7 days after placing in the greenhouse).